

Applicants : Qiwang Xu et al.
Appln. No. : 10/524,476
Page : 2

In the Claims:

This listing of claims will replace all prior versions and listings of claims in the application:

1. (previously presented) A method for treating a subject suffering from urogenital tract infection, the method comprising administering to the subject an effective amount of a medicament containing N-acetyl-D-glucosamine and/or pharmaceutical acceptable salts thereof.
2. (previously presented) The method according to claim 1, wherein said medicament is a formulation in the form for topical administration.
3. (previously presented) The method according to claim 2, wherein said medicament is a formulation in a form selected from a group consisting of lotion, aqua, emulsion, cream, ointment, and suppository.
4. (previously presented) The method according to claim 1, wherein a daily dose of the administration is 100 to 2000 mg of N-acetyl-D-glucosamine.
5. (previously presented) The method according to claim 1, wherein a concentration of N-acetyl-D-glucosamine in said medicament is 0.1 to 10% by weight.
6. (previously presented) The method according to claim 2, wherein a daily dose of the administration is 100 to 2000 mg of N-acetyl-D-glucosamine.
7. (previously presented) The method according to claim 3, wherein a daily dose of the administration is 100 to 2000 mg of N-acetyl-D-glucosamine.
8. (previously presented) The method according to claim 2, wherein a concentration of N-acetyl-D-glucosamine in said medicament is 0.1 to 10% by weight.

Applicants : Qiwang Xu et al.
Appln. No. : 10/524,476
Page : 3

9. (previously presented) The method according to claim 3, wherein a concentration of N-acetyl-D-glucosamine in said medicament is 0.1 to 10% by weight.

10. (new) A method for treating a subject suffering from urogenital tract infection, the method comprising administering to the subject an effective amount of a medicament containing as the only active ingredient N-acetyl-D-glucosamine and/or pharmaceutically acceptable salts thereof.

11. (new) The method according to claim 10, wherein said medicament is a formulation in the form for topical administration.

12. (new) The method according to claim 11, wherein said medicament is a formulation in a form selected from a group consisting of lotion, aqua, emulsion, cream, ointment, and suppository.

13. (new) The method according to claim 10, wherein a daily dose of the administration is 100 to 2000 mg of N-acetyl-D-glucosamine.

14. (new) The method according to claim 10, wherein a concentration of N-acetyl-D-glucosamine in said medicament is 0.1 to 10% by weight.

15. (new) The method according to claim 11, wherein a daily dose of the administration is 100 to 2000 mg of N-acetyl-D-glucosamine.

16. (new) The method according to claim 12, wherein a daily dose of the administration is 100 to 2000 mg of N-acetyl-D-glucosamine.

17. (new) The method according to claim 11, wherein a concentration of N-acetyl-D-glucosamine in said medicament is 0.1 to 10% by weight.

Applicants : Qiwang Xu et al.
Appln. No. : 10/524,476
Page : 4

18. (new) The method according to claim 12, wherein a concentration of N-acetyl-D-glucosamine in said medicament is 0.1 to 10% by weight.